





ORIGINAL ARTICLE

A novel strain of lumpy skin disease virus causes clinical disease in cattle in Hong Kong

John Flannery¹  | Barbara Shih²  | Ismar R. Haga¹ | Martin Ashby¹ | Amanda Corla¹ | Simon King¹  | Graham Freimanis¹ | Noemi Polo¹ | Anne Ching-nga Tse³ | Christopher J. Brackman³ | Jason Chan³ | Patrick Pun³ | Andrew D. Ferguson^{3,4} | Andy Law²  | Samantha Lycett² | Carrie Batten¹ | Philippa M. Beard^{1,2}

¹ The Pirbright Institute, Woking, Surrey, UK

² The Roslin Institute, University of Edinburgh, Midlothian, UK

³ Agriculture, Fisheries and Conservation Department, Government of the Hong Kong Special Administrative Region, Hong Kong, China

⁴ CityU Veterinary Diagnostic Laboratory, City University of Hong Kong, Hong Kong, China

Correspondence

Philippa M. Beard, The Pirbright Institute, Ash Rd, Surrey, UK.

Email: pip.beard@pirbright.ac.uk

John Flannery, Barbara Shih, and Ismar R. Haga are the joint first authors.

Samantha Lycett, Carrie Batten, and Philippa M. Beard are the joint senior authors.

Funding information

Rural and Environment Science and Analytical Services Division; Department for Environment, Food and Rural Affairs, UK Government; Biotechnology and Biological Sciences Research Council

Abstract

Lumpy skin disease virus (LSDV) is an emerging poxviral pathogen of cattle that is currently spreading throughout Asia. The disease situation is of high importance for farmers and policy makers in Asia. In October 2020, feral cattle in Hong Kong developed multi-focal cutaneous nodules consistent with lumpy skin disease (LSD). Gross and histological pathology further supported the diagnosis and samples were sent to the OIE Reference Laboratory at The Pirbright Institute for confirmatory testing. LSDV was detected using quantitative polymerase chain reaction (qPCR) and additional molecular analyses. This is the first report of LSD in Hong Kong. Whole genome sequencing (WGS) of the strain LSDV/Hong Kong/2020 and phylogenetic analysis were carried out in order to identify connections to previous outbreaks of LSD, and better understand the drivers of LSDV emergence. Analysis of the 90 core poxvirus genes revealed LSDV/Hong Kong/2020 was a novel strain most closely related to the live-attenuated Neethling vaccine strains of LSDV and more distantly related to wildtype LSDV isolates from Africa, the Middle East and Europe. Analysis of the more variable regions located towards the termini of the poxvirus genome revealed genes in LSDV/Hong Kong/2020 with different patterns of grouping when compared to previously published wildtype and vaccine strains of LSDV. This work reveals that the LSD outbreak in Hong Kong in 2020 was caused by a different strain of LSDV than the LSD epidemic in the Middle East and Europe in 2015–2018. The use of WGS is highly recommended when investigating LSDV disease outbreaks.

KEYWORDS

cattle, epidemic, Hong Kong, Lumpy skin disease, lumpy skin disease virus, phylogenetics, poxvirus

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Transboundary and Emerging Diseases* published by Wiley-VCH GmbH

1 | INTRODUCTION

Lumpy skin disease (LSD) is a severe disease of cattle and water buffalo characterized by multi-focal cutaneous nodules. It is caused by infection with the poxvirus lumpy skin disease virus (LSDV), a member of the capripoxvirus genus. LSDV is transmitted by hematophagous vectors, such as mosquitoes and flies, which facilitate rapid spread of the virus in optimal climatic conditions (Sanz-Bernardo et al., 2021). LSDV is a rapidly emerging pathogen, having spread over the past ten years from Africa and the Middle East into south-east Europe, the Caucasus, Russia and, more recently, Asia (Acharya & Subedi, 2020; Beard, 2016; Calistri et al., 2020; Kumar et al., 2021; Lu et al., 2020; Sudhakar et al., 2020). This Eurasian LSD epidemic has affected thousands of cattle and caused substantial economic loss through the loss of animals, reduced productivity, the cost of control and prevention campaigns, and loss of export markets (Casal et al., 2018). Widespread vaccination programmes with the live-attenuated LSDV vaccine based on the Neethling strain of the virus have been a key to the control of the disease in south-east Europe (Efsa, 2018; Klement et al., 2018).

Phylogenetic analysis of LSDV isolates using whole genome sequencing (WGS) has revealed that strains sequenced to date segregate into two subgroups (Biswas et al., 2020). Subgroup 1.1 contains vaccine strains related to the original Neethling strain. Subgroup 1.2 can be divided into wildtype LSDV strains and LSDV KSGP strains. Examination of wildtype LSDV strains shows that the viral strains which caused disease in south-east Europe in 2015–2016 and Russia in 2015 are very similar to a strain isolated from Israel in 2012, supporting the theory of northward spread of the virus from Africa and the Middle East. In contrast, WGS of a novel strain of LSDV, known as Saratov/2017, from diseased cattle in southern Russia in 2017 revealed similarities with the Neethling vaccine strain (Sprygin et al., 2018). The origin of Saratov/2017 is unclear.

We report here the clinical and pathological features of the first reported outbreak of LSD in Hong Kong in 2020, and phylogenetic analysis of the whole genome of the LSDV strain isolated from affected cattle (LSDV/HongKong/2020). This strain differs from all previously reported LSDV strains but clusters phylogenetically with Neethling vaccine strains.

2 | MATERIAL AND METHODS

Tissue samples for histopathological analysis were fixed in formalin, processed to paraffin wax blocks, sectioned, and stained with haematoxylin and eosin. EDTA blood samples ($n = 2$), tissues ($n = 9$), nasal swab ($n = 1$), and serum ($n = 2$) from three cattles were submitted to the OIE reference laboratory for LSD at The Pirbright Institute, UK, for confirmatory diagnosis. Nucleic acid was extracted from EDTA blood samples, tissues, and swabs using the MagMAX™ CORE Nucleic Acid Purification Kit (ThermoFisher Scientific, Paisley, UK) on the KingFisher Flex extraction platform (ThermoFisher Scientific). Capripoxvirus DNA was detected by a qPCR assay targeting LSDV074

(also known as the p32 gene, a homolog of the vaccinia virus H3L gene) (Bowden et al., 2008) using the Path-ID™ qPCR Master Mix (ThermoFisher Scientific). Nucleic acid from capripoxvirus positive samples was further analyzed by a qPCR targeting LSDV011 designed to differentiate the three capripoxvirus species (Lamien et al., 2011) and a qPCR targeting LSDV008 designed to differentiate LSDV wildtype strains from Neethling vaccine strains (Sprygin et al., 2018). Partial genome sequencing of LSDV036 (the RNA polymerase subunit RPO30) was performed using primers described by Gelaye et al. (2015) and the resulting sequences were assembled using the SeqMan Pro software and phylogenetic analysis performed using MEGA7. Serum samples were tested with the ID Screen® Capripox Double-antigen ELISA (ID Vet Innovative diagnostics) following the manufacturer's instructions.

LSDV was isolated on MDBK cells (ATCC code CCL-22). Briefly, tissue homogenates from fresh skin were prepared from each of the three cattles, homogenates were sonicated twice and centrifuged prior to inoculation onto MDBK cell monolayers. Cells were harvested after 6 days and then sonicated prior to a second passage on MDBK cells using both cellular and supernatant fractions. The cells were harvested when 100% CPE was observed.

LSDV was purified as described previously (Fay et al., 2020). Briefly, MDBK cells infected with LSDV were harvested, centrifuged at low speed, supernatants discarded, and the pellets resuspended in 1 mM of Tris-HCl pH9. Samples were vortexed, sonicated, and incubated with Benzonase® (>250 units/μL, Sigma E1014-25KU) ahead of two rounds of sucrose cushion purification. Samples were then treated with 33 μL of 1.5 M Tris pH 8.8, 50 μL of 10% SDS, 100 μL of 60% sucrose, and 85 μL of proteinase K (20 mg/mL, ThermoFisher Scientific) for 4 h at 37°C, followed by phenol-chloroform extraction and ethanol precipitation.

Total dsDNA (1 ng) was processed for WGS using the Nextera XT DNA library kit (Illumina) and performed on the Hamilton NGSStar (Hamilton Robotics). Libraries were bead normalized and pooled according to manufacturer's protocols. Final sequencing pools were loaded at 12 pM concentration onto an Illumina MiSeq 300 cycle sequencing run with a 1% PhiX spike-in (Illumina).

Using BBtools (Bushnell), adaptor and quality trimming were carried out on the reads before genome assembly. Two approaches were taken; the consensus sequence and the de novo assembly methods. For generating the consensus sequence, reads were mapped to the LSDV isolate 155920/2012 (KX894508.1) with BWA-MEM (version 0.7.17) (Li & Durbin, 2009), and SAMtools (version 1.10) (Li et al., 2009) was used for processing the alignment files. The variants were detected using freebayes (version 1.3.1) (Garrison & Marth, 2012), and the per-region coverage was estimated using mosdepth (version 0.2.6). BCFtools (version 1.10) was used for creating the consensus sequence using high-quality calls (%QUAL ≥ 20), taking into account the regions with zero coverage from the mosdepth output. De novo assembly was carried out using Spades (version 3.13.2) (Prjibelski et al., 2020), setting the kmers to "33, 55, 77, 99." Using RaGOO (Alonge et al., 2019), reference-guided scaffolding was carried out on the contigs generated by SPAdes with the consensus sequence acting as the template. The consensus and de novo

genome sequences were compared to each other using Minimap2 (version 2.16) (Li, 2018). Custom Python scripts were used to combine the two genome sequences into the final LSDV/HongKong/2020 genome sequence; in brief, where there are differences in variants, the variants in the *de novo* assembly were preferentially included, and stretches of poly-*N* were retained only if it was present in both the consensus and *de novo* sequences.

Prokka (Seemann, 2014) was used for annotating the LSDV/HongKong/2020 genome, along with 21 other lumpy skin disease genomes on NCBI (MH646674.1, AF409138.1, KX764645.1, KX764644.1, MG972412.1, KX764643.1, MT134042.1, MN072619.1, AF325528.1, KX683219.1, AF409137.1, MH893760.2, KY702007.1, MN642592.1, KX894508.1, KY829023.3, MN995838.1, MT007950.1, MT643825.1, MT992618.1, NC_003027.1, see Table 1 for details). An all-against-all comparison was made on the Prokka-identified gene sequences using BLAST+ (version 2.7.1) (Camacho et al., 2009). Network analysis based on the BLAST output was performed using Graphia (Freeman et al., 2020) for labelling the Prokka-identified genes with corresponding genes in the reference genome annotation (KX894508.1) in order to find a set of core genes between the LSDV genomes and to find groups of accessory genes. Each sequence (i.e., a gene in a genome) was treated as a node, two sequences were joined by an edge if they had significant similarity score (≥ 60 bitscore, ≥ 60 qcovs, and $evalue < 1e-5$). Markov Clustering was used for grouping nodes sharing high similarity into clusters, with each cluster containing sequences of the same gene from different genomes. In order to highlight genes that showed a strong difference between vaccine and wildtype strains, we explored the accessory gene clusters that separated to distinct components when applying an additional edge filter using 14 kNN (k-nearest neighbour) based on bitscore.

For each of the 22 genomes, the core gene sequences were merged into one continuous sequence for building a phylogenetic tree (for consecutive core genes, the intergenic regions between the genes were also included). MUSCLE (MULTiple Sequence Comparison by Log-Expectation; version 3.8.1551) (Madeira et al., 2019) was used for aligning the core gene sequences for the 22 genomes. RAXML (Randomized Axelerated Maximum Likelihood; version 8.2.12) was used to build the phylogenetic tree using a general time reversible substitution model with gamma-distributed site to site rate variation (GTRGAMMA) and 100 bootstraps, which was then visualized using iTOL (Letunic & Bork, 2019).

3 | RESULTS AND DISCUSSION

There are no commercial cattle farms in Hong Kong, however, there is a population of around 1000 feral brown cattle found mostly in local country parks. There is also a population of some 200 feral water buffaloes. In October 2020, multiple brown cattle in the eastern part of the New Territories of Hong Kong were reported to have multi-focal nodular skin lesions (Figure 1A and B). In early November 2020, similar cases were detected in the northern New Territories close to the bor-

der with mainland China and on one of the outlying islands in the southwest. In addition to the skin lesions, other clinical signs included fever, malaise, anorexia, and superficial lymphadenopathy. Nasal and/or oral ulcers were present in some cattle (Figure 1C), accompanied by nasal discharges and/or ptyalism. The clinical course lasted for 2–3 weeks and the disease was self-limiting in a majority of cases. The morbidity of the initial outbreak based on clinical signs was estimated to be between 20 and 30%. Two affected cattles were reported dead, however, both were over 15 years old with only mild skin lesions. It was uncertain whether death was causally associated with LSD or other unknown comorbidities. Clinical cases were absent from the buffalo populations.

Postmortem examination was carried out on two cattles at the Government's official veterinary laboratory, Tai Lung Veterinary Laboratory. Gross and microscopic findings were consistent with a diagnosis of LSD (Sanz-Bernardo et al., 2020). Gross findings included widespread, randomly distributed cutaneous, and subcutaneous nodules ranging from 1 to 40 mm in diameter, sometimes with a targetoid appearance. A few nodules were ulcerated and others had a central area of dense crust (*sit-fast*). Multiple superficial lymph nodes were enlarged and haemorrhagic, especially the prescapular lymph nodes. The most striking microscopic lesion was a necrotizing vasculitis that often started from the deep cutaneous plexus with abundant surrounding infiltrates of predominate large histiocytes and fibroblasts (Figure 2A). The histiocytes frequently contained a large, prominent eosinophilic or amphophilic, intracytoplasmic inclusion bodies and had marginated chromatin (Figure 2B). Other lesions included epidermal ballooning degeneration, sub-corneal vesicles, and intracytoplasmic inclusion bodies within keratinocytes.

The clinical features of LSD caused by LSDV/HongKong/2020 strain in the field appear similar to those reported from outbreaks in south-east Europe, Africa, and Asia. Previous studies have reported morbidities from 1 to 20% and mortality of 1% or less (Acharya & Subedi, 2020; European Food Safety, 2018; Kumar et al., 2021; Lu et al., 2020; Sudhakar et al., 2020) which are consistent with the morbidity and mortality estimates of 20–30% and 0% reported in the outbreak in Hong Kong. Challenge studies carried out under experimental conditions will be required to compare in more detail the virulence of LSDV/HongKong/2020 with wildtype and KSGP strains.

Tissue samples, including subcutaneous nodules and enlarged lymph nodes, were submitted to the OIE Reference Laboratory for LSD at The Pirbright Institute, UK. The capripoxvirus gene LSDV074 was detected in EDTA blood, tissue, and swab samples ($n = 12$), confirming the clinical diagnosis. LSDV was then speciated in each sample using the differentiation assay targeting LSDV011. Finally, LSDV Neethling vaccine strain was identified using the vaccine-specific assay targeting LSDV008. Both serum samples were positive for capripoxvirus antibodies. The RP030 full gene sequences were amplified and compared to capripoxvirus reference strains and identified as LSDV with the closest relatives being Neethling vaccine strains (AF409138, KX764643, KX764644, KX764645).

LSDV was isolated from skin samples taken from all three animals and named LSDV/HongKong/2020/01 to 03. Viral DNA

TABLE 1 Sequences included in the analysis

| GenBank ID | Abbreviated name | Year | Origin | Wildtype or vaccine | Identity to HongKong/2020 (MW732649) | Identity to vaccine (AF409138) | Identity to wildtype (KX894508) |
|-------------|------------------------|------|--------------------|----------------------|--------------------------------------|--------------------------------|---------------------------------|
| AF32528.1 | Ni-2490 | 1959 | Kenya | wildtype | 99.28 | 98.77 | 99.89 |
| AF409137.1 | Neethling Warmbaths LW | 1999 | South Africa | wildtype | 99.26 | 98.78 | 99.97 |
| KX894508.1 | Israel/2012 | 2012 | Israel | wildtype | 99.26 | 98.78 | 100 |
| KY702007.1 | Bunjanovac/2016 | 2016 | Bunjanovac, Serbia | wildtype | 99.27 | 98.78 | 100 |
| KY829023.3 | Evros/2015 | 2015 | Evros, Greece | wildtype | 99.27 | 98.78 | 100 |
| MH646674.1 | Saratov/2017 | 2017 | Saratov, Russia | wildtype | 99.44 | 99.53 | 99.19 |
| MH893760.2 | Dagestan/2015 | 2015 | Dagestan, Russia | wildtype | 99.23 | 98.76 | 99.96 |
| MN072619.1 | Kenya/1958 | 1958 | Kenya | wildtype | 99.3 | 98.77 | 99.91 |
| MN642592.1 | Kubash/2016 | 2016 | Kubash, Kazakhstan | wildtype | 99.27 | 98.78 | 100 |
| MT134042.1 | Udmurtiya/2019 | 2019 | Udmurtiya, Russia | wildtype | 99.44 | 99.31 | 99.37 |
| MW732649.1 | HongKong/2020 | 2020 | Hong Kong, China | wildtype | 100 | 99.42 | 99.26 |
| MN995838.1 | Turkey/2014 | 2014 | Turkey | wildtype | 99.26 | 98.78 | 99.99 |
| MT007950.1 | Namibia/2016 | 2016 | Namibia | wildtype | 99.18 | 98.69 | 99.89 |
| MT643825.1 | Bulgaria/2016 | 2016 | Bulgaria | wildtype | 99.26 | 98.78 | 100 |
| MT992618.1 | Kazakhstan/2018 | 2018 | Kazakhstan | wildtype | 99.41 | 99.34 | 99.39 |
| NC_003027.1 | Ni-2490 | | | wildtype | 99.28 | 98.77 | 99.89 |
| AF409138.1 | Neethling LW | | | Vaccine | 99.42 | 100 | 98.78 |
| KX683219.1 | KSGP 0240 | | | Vaccine ^a | 99.3 | 98.77 | 99.91 |
| KX764643.1 | Neethling SIS | | | Vaccine | 99.42 | 100 | 98.77 |
| KX764644.1 | Neethling Herbivac | | | Vaccine | 99.42 | 100 | 98.77 |
| KX764645.1 | Neethling OBP | | | Vaccine | 99.42 | 100 | 98.78 |
| MG972412.1 | Neethling Croatia | | | Vaccine | 99.42 | 99.99 | 98.77 |

^aHas been discontinued as a live attenuated vaccine against LSDV. The percentage similarity was calculated by accounting only aligned positions where both the query and subject genomes have sequences, thus ignoring insertions, deletions and regions without coverage.



FIGURE 1 Clinical presentation of LSD in a feral brown bull, Hong Kong, 2020. (A, B) Multi-focal, raised skin nodules. (C) Focal ulceration of the nasal mucosa

from LSDV/HongKong/2020/01 was purified, sequenced, and the genome assembled. The genome sequence has been named LSDV/HongKong/2020 and deposited in Genbank [MW732649.1]. The genome of LSDV/HongKong/2020 was compared to published genomes of other LSDV strains (Table 1). Comparison of the 90 core genes that are conserved in all chordopoxviruses (Gubser et al., 2004) revealed that LSDV/HongKong/2020 was most closely related to Neethling strains of LSDV and the Saratov/2017 strain, and more distantly related to LSDV isolates from the Middle East, Europe, and neighbouring regions (Figure 3). This reveals that the LSD outbreak in Hong Kong in 2020 and the LSD outbreaks in south-east Europe in 2015–2017 were caused by different strains of LSDV.

More detailed analysis of the accessory genes of the LSDV genomes revealed 29 genes that formed separate groupings in the network analysis when a more stringent kNN filter was applied. In Figure 3, those

that grouped with the vaccine strain Neethling LW are labelled red, and those that grouped with the wildtype LSDV strain 155920/2012 are labelled blue. Six different patterns of grouping across the LSDV genomes were noted (a-f). Regardless of the genes, most wildtype and KSGP isolates grouped with wildtype LSDV strain 155920/2012, whereas the Neethling strains grouped with Neethling LW. However, four LSDV strains each showed unique patterns of these accessory gene alleles—Saratov/2017, Udmurtiya/2019, Kazakhstan/2018, and LSDV/HongKong/2020. These four strains shared higher similarity to the wildtype LSDV strain 155920/2012 for some accessory genes and higher similarity to the Neethling LW vaccine strain for other accessory genes.

The implications of the identification of the novel LSDV/HongKong/2020 strain on the diagnosis and control of LSD were considered. Diagnostic qPCR assays to differentiate between




FIGURE 2 LSD causes marked, multi-focal, necrotizing, and histiocytic dermatitis with (A) necrotizing vasculitis and (B) intracytoplasmic inclusion bodies

infected and vaccinated animals should be used with caution, as in this instance we were unable to use the LSDV vaccine-like assay (Sprygin et al., 2018) to differentiate between LSDV/HongKong/2020 and the Neethling vaccine strain due to sequence similarity within the amplified region. Redesigning these assays will be required to enable differentiation of this newly circulating strain from Neethling strain vaccines. Currently available live-attenuated LSDV vaccines are likely to provide protection against LSDV/HongKong/2020 since poxviruses are known to provide broad within-genus protection, however, vaccination and challenge studies should be undertaken to confirm this.

This study has shown the value of sequencing the whole genome of LSDV rather than a small number of individual genes. We recommend carrying out WGS of new isolates of LSDV whenever possible and making the data available promptly to allow rapid identification of new LSDV strains and facilitate understanding of LSDV epidemics.

Note added in proof

While this manuscript was under review a manuscript describing a LSDV outbreak in the southeast China in 2020 was released




FIGURE 3 Core gene phylogenetic tree. The phylogenetic tree was built using the core genes (including intergenic regions between consecutive core genes) using RAXML and midpoint rooted. For accessory genes, isolates that grouped with either vaccine strains or wildtype strains in the network analysis are highlighted with red and blue respectively (panel on right). Different patterns of grouping are seen for 29 accessory genes: (A) LD141, LD142, LD146, LD150, (B) LD015, LD017, LD035, LD127, LD151, (C) LD006, LD143, (D) LD009, LD135, LD137, LD138, (E) LD008, LD010, (F) LD018, LD020, LD021, LD023, LD066, LD067, LD122, LD125, LD126, LD128, LD129, LD130

(doi.org/10.1101/2021.04.27.441643), including the full genome sequence of the LSDV strain from this outbreak. Initial comparisons indicate this strain is very closely related to the HongKong/2020 strain described in this manuscript.

ACKNOWLEDGMENTS

The Pirbright Institute is funded by the Department for Environment, Food and Rural Affairs (Grant code: SE26081) and the Biotechnology and Biological Sciences Research Council (BBSRC) (Grant code: BB/R008833/1). IRH and PMB were supported by BBSRC strategic funding to the Pirbright Institute (Grant codes: BBS/E/I/00007036, BBS/E/I/00007037, and BBS/E/I/00007039). BS, AL, SL, and PMB are supported by a BBSRC Institute Strategic Programme grant to Roslin Institute (Grant code: BBS/E/D/20002173), and SL is additionally supported by Scottish Government Rural and Environment Science and Analytical Services Division as part of Centre of Expertise on Animal Disease Outbreaks (EPIC).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ETHICAL GUIDELINES

The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines page, have been adhered to. No ethical approval was required for this study.

AUTHOR CONTRIBUTIONS

J.F., B.S., I.R.H., A.L., S.L., C.B., and P.M.B. conceptualised the study; J.C. carried out the clinical analyses; B.S., I.R.H., M.A., A.C., G.F., S.K., A.T., C.J.B., J.C., P.P., N.P. carried out the laboratory assays; A.D.F. carried out the necropsy and histopathology; J.F., S.K., B.S., I.R.H., A.L., S.L., and P.M.B. analysed the data; P.M.B. drafted the paper. All authors discussed the results, commented on the manuscript, and gave final approval of the version to be published.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in Genbank at <https://www.ncbi.nlm.nih.gov/genbank/>; reference number MW732649.1.

ORCID

John Flannery <https://orcid.org/0000-0002-5012-2829>

Barbara Shih <https://orcid.org/0000-0002-3676-3304>

Simon King <https://orcid.org/0000-0002-3129-2961>

Andy Law <https://orcid.org/0000-0003-1868-2364>

REFERENCES

- Acharya, K. P., & Subedi, D. (2020). First outbreak of lumpy skin disease in Nepal. *Transbound Emerg Dis*, 67(6), 2280-2281.
- Alonge, M., Soyk, S., Ramakrishnan, S., Wang, X., Goodwin, S., Sedlazeck, F. J., Lippman, Z. B., & Schatz, M. C. (2019). RaGOO: Fast and accurate reference-guided scaffolding of draft genomes. *Genome Biology*, 20(1), 224.

- Beard, P. M. (2016). Lumpy skin disease: A direct threat to Europe. *Veterinary Record*, 178(22), 557-558.
- Biswas, S., Noyce, R. S., Babiuk, L. A., Lung, O., Bulach, D. M., Bowden, T. R., Boyle, D. B., Babiuk, S., & Evans, D. H. (2020). Extended sequencing of vaccine and wild-type capripoxvirus isolates provides insights into genes modulating virulence and host range. *Transbound Emerg Dis*, 67(1), 80-97.
- Bowden, T. R., Babiuk, S. L., Parkyn, G. R., Copps, J. S., & Boyle, D. B. (2008). Boyle DBCapripoxvirus tissue tropism and shedding: A quantitative study in experimentally infected sheep and goats. *Virology*, 371(2), 380-393.
- Bushnell, B. BMap. Retrieved from: sourceforge.net/projects/bbmap/.
- Calistri, P., De Clercq, K., Gubbins, S., Klement, E., Stegeman, A., Cortiñas Abrahantes, J., Marojevic, D., Antoniou, S. E., & Broglia, A. (2020). Lumpy skin disease epidemiological report IV: Data collection and analysis. *Efsa Journal*, 18(2), e06010.
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: Architecture and applications. *Bmc Bioinformatics*, 10, 421.
- Casal, J., Allepuz, A., Miteva, A., Pite, L., Tabakovsky, B., Terzievski, D., Alexandrov, T., & Beltrán-Alcrudo, D. (2018). Economic cost of lumpy skin disease outbreaks in three Balkan countries: Albania, Bulgaria and the Former Yugoslav Republic of Macedonia (2016-2017). *Transbound Emerg Dis*, 65(6), 1680-1688.
- Efsa. (2018). Lumpy skin disease II. Data collection and analysis. *Efsa Journal*, 16(2), e05176.
- European Food Safety Authority. (2018). Lumpy skin disease II. Data collection and analysis. *EFSA J*, 16(2), e05176.
- Fay, P. C., Cook, C. G., Wijesiriwardana, N., Tore, G., Comtet, L., Carpentier, A., Shih, B., Freimanis, G., Haga, I. R., & Beard, P. M. (2020). Madin-Darby bovine kidney (MDBK) cells are a suitable cell line for the propagation and study of the bovine poxvirus lumpy skin disease virus. *Journal of Virological Methods*, 285, 113943.
- Freeman, T. C., Horsewell, S., Patir, A., Harling-Lee, J., Regan, T., Shih, B. B., Prendergast, J., Hume, D. A., & Angus, T. (2020). Graphia: A platform for the graph-based visualisation and analysis of complex data. *bioRxiv*, 2020.09.02.279349.
- Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read sequencing. *arXiv preprint arXiv:1207.3907*.
- Gelaye, E., Belay, A., Ayelet, G., Jenberie, S., Yami, M., Loitsch, A., Tuppurainen, E., Grabherr, R., Diallo, A., & Lamien, C. E. (2015). Capripox disease in Ethiopia: Genetic differences between field isolates and vaccine strain, and implications for vaccination failure. *Antiviral Research*, 119, 28-35.
- Gubser, C., Hué, S., Kellam, P., & Smith, G. L. (2004). Poxvirus genomes: A phylogenetic analysis. *Journal of General Virology*, 85(Pt 1), 105-117.
- Klement, E., Broglia, A., Antoniou, S. E., Tsiamadis, V., Plevraki, E., Petrović, T., Polaček, V., Debeljak, Z., Miteva, A., Alexandrov, T., Marojevic, D., Pite, L., Kondratenko, V., Atanasov, Z., Gubbins, S., Stegeman, A., & Abrahantes, J. C. (2018). Neethling vaccine proved highly effective in controlling lumpy skin disease epidemics in the Balkans. *Preventive Veterinary Medicine*, 181, 104595.
- Kumar, N., Chander, Y., Kumar, R., Khandelwal, N., Riyesh, T., Chaudhary, K., Shanmugasundaram, K., Kumar, S., Kumar, A., Gupta, M. K., Pal, Y., Barua, S., & Tripathi, B. N. (2021). Isolation and characterization of lumpy skin disease virus from cattle in India. *Plos One*, 16(1), e0241022.
- Lamien, C. E., Lelenta, M., Goger, W., Silber, R., Tuppurainen, E., Matijevic, M., Luckins, A. G., & Diallo, A. (2011). Real time PCR method for simultaneous detection, quantitation and differentiation of capripoxviruses. *Journal of Virological Methods*, 171(1), 134-140.
- Letunic, I., & Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: Recent updates and new developments. *Nucleic Acids Research*, 47(W1), W256-W259.
- Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*, 34(18), 3094-3100.
- Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, 25(14), 1754-1760.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin, R.; 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16), 2078-2079.
- Lu, G., Xie, J., Luo, J., Shao, R., Jia, K., & Li, S. (2020). Lumpy skin disease outbreaks in China, since 3 August 2019. *Transbound Emerg Dis*, 68(2), 216-219.
- Madeira, F., Park, Y. M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey, A. R. N., Potter, S. C., Finn, R. D., & Lopez, R. (2019). Lopez RThe EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Research*, 47(W1), W636-W641.
- Prijbelski, A., Antipov, D., Meleshko, D., Lapidus, A., & Korobeynikov, A. (2020). Using SPAdes de novo assembler. *Current Protocols in Bioinformatics*, 70(1), e102.
- Sanz-Bernardo, B., Haga, I. R., Wijesiriwardana, N., Basu, S., Larner, W., Diaz, A. V., Langlands, Z., Denison, E., Stoner, J., White, M., Sanders, C., Hawes, P. C., Wilson, A. J., Atkinson, J., Batten, C., Alphey, L., Darpel, K. E., Gubbins, S., & Beard, P. M. (2021). Quantifying and modelling the acquisition and retention of lumpy skin disease virus by haematophagus insects reveals clinically but not subclinically-affected cattle are promoters of viral transmission and key targets for control of disease outbreaks. *Journal of Virology*, 95(9), e02239-20.
- Sanz-Bernardo, B., Haga, I. R., Wijesiriwardana, N., Hawes, P. C., Simpson, J., Morrison, L. R., Macintyre, N., Brocchi, E., Atkinson, J., Haegeman, A., De Clercq, K., Darpel, K. E., & Beard, P. M. (2020). Beard PMLumpy skin disease is characterized by severe multifocal dermatitis with necrotizing fibrinoid vasculitis following experimental infection. *Veterinary Pathology*, 57(3), 388-396.
- Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*, 30(14), 2068-2069.
- Sprygin, A., Babin, Y., Pestova, Y., Kononova, S., Wallace, D. B., Van Schalkwyk, A., Byadovskaya, O., Diev, V., Lozovoy, D., & Kononov, A. (2018). Analysis and insights into recombination signals in lumpy skin disease virus recovered in the field. *Plos One*, 13(12), e0207480.
- Sprygin, A., Pestova, Y., Prutnikov, P., & Kononov, A. (2018). Detection of vaccine-like lumpy skin disease virus in cattle and *Musca domestica* L. flies in an outbreak of lumpy skin disease in Russia in 2017. *Transbound Emerg Dis*, 65(5), 1137-1144.
- Sudhakar, S. B., Mishra, N., Kalaiyarasu, S., Jhade, S. K., Hemadri, D., Sood, R., Bal, G. C., Nayak, M. K., Pradhan, S. K., & Singh, V. P. (2020). Lumpy skin disease (LSD) outbreaks in cattle in Odisha state, India in August 2019: Epidemiological features and molecular studies. *Transbound Emerg Dis*, 67(6), 2408-2422.

How to cite this article: Flannery, J., Shih, B., Haga, I. R., Ashby, M., Corla, A., King, S., Freimanis, G., Polo, N., Tse, A. C.-n., Brackman, C. J., Chan, J., Pun, P., Ferguson, A. D., Law, A., Lycett, S., Batten, C., & Beard, P. M. (2021). A novel strain of lumpy skin disease virus causes clinical disease in cattle in Hong Kong. *Transboundary and Emerging Diseases*, 1-8. <https://doi.org/10.1111/tbed.14304>